

Problem set 2 (due October 27)

- Consider a point particle lying a distance $h \ll R$ from the surface of the Earth, the latter of which is assumed to be a sphere of radius R rotating with constant angular velocity w . The particle is subject to the force of gravity and starts at rest at a point $p_0 = (r_0, \varphi_0, z_0) = (R + h, 0, 0)$ in cylindrical coordinates, i.e. it starts a distance h above the equator with velocity $\vec{v} = 0$. With this assumption, the particle is confined to move on the $z = 0$ plane.
 - (1pt) How many degrees of freedom are needed to describe the motion of the particle?
 - (2pt) Find the Lagrangian, identify the cyclic coordinates, and the corresponding conserved quantities
 - (2pt) Using the initial conditions and conservation laws, write down the Euler-Lagrange equations in terms of w and r_0
 - (2pt) Assuming that $\left(\frac{wr_0^2}{r^2}\right)^2 \ll \frac{g}{r}$ solve the equations of motion and find how much the particle deviates from φ_0 when it reaches the ground
- In class, we showed that for every continuous symmetry of the Lagrangian there exists a conserved quantity. Consider now the more general family of maps

$$q_i(t) \mapsto Q_i(s, t), \quad t \mapsto T(s, t) \quad \text{where} \quad Q_i(0, t) = q_i(t), \quad T(0, t) = t \quad (1)$$

where $s \in \mathbb{R}$ and

$$\left. \frac{\partial^2 T}{\partial s \partial t} \right|_{s=0} = \left. \frac{d}{dt} \frac{\partial T}{\partial s} \right|_{s=0} \quad (2)$$

This family of maps is a continuous symmetry of the Lagrangian whenever the following one-form is independent of s ,

$$L(Q_i(s, t), \dot{Q}_i(s, t), T) dT(s, t) + \frac{d}{dT} F(Q_i(s, t), T(s, t), s) dT(s, t) \quad (3)$$

where $F(Q_i(s, t), T(s, t), 0) = 0$. In particular, note that

$$\dot{Q} = \frac{dQ}{dT} = \frac{\partial T}{\partial t} \left(\frac{\partial Q_i}{\partial q_j} \frac{\partial q_j}{\partial t} + \frac{\partial Q_i}{\partial t} \right) \quad (4)$$

- (3pts) Find the conserved quantity $J(q_i, \dot{q}_i, t)$ associated with this symmetry (hints: (1) you can extract J from an expression of the form $(\frac{d}{dt} J) dt = 0$; (2) you will need to show and use $\frac{\partial L}{\partial t} = -\frac{d}{dt} (\dot{q}_i \frac{\partial L}{\partial \dot{q}_i} - L)$)

- (2pts) Show that the following action is invariant under all time reparametrizations $t \mapsto T(t)$

$$S = \int \sqrt{\dot{\vec{r}} \cdot \dot{\vec{r}}} dt \quad (5)$$

Letting $T = t + s_n t^n$, where s_n is a constant, compute the conserved quantity associated with each s_n .

Bonus (1pt): what's the interpretation of this result?


(c) (2pts) Consider a system of particles with Lagrangian

$$L = \frac{1}{2} m_i |\dot{\vec{r}}_i|^2 - V(r_i - r_j) \quad (6)$$

Identify the function F and compute the conserved current associated with a (Galilean) boost $\vec{r}_i \rightarrow \vec{r}_i + \vec{v}t$. This completes the derivation of conserved quantities associated with Galilean transformations discussed in class.

Bonus (1pt): what's the physical interpretation to this result?

3. Consider a pendulum that consists of a point particle of mass m hanging from a block of mass M by a massless rod of length l . The block moves in the horizontal direction under the influence of a potential $V(x)$, as illustrated below



- (a) (1pts) How many degrees of freedom are needed to describe the evolution of the system?
- (b) (2pts) Write down the Lagrangian and the corresponding equations of motion
- (c) (2pts) Let $V(x) = kx^2$ and obtain an ordinary second-order differential equation for the motion of m
- (d) (1pt) Solve the equations of motion in the small angle approximation.

Bonus (1pt): find the choice of parameters for which the answer diverges and the small angle approximation breaks down. What's the physical interpretation of this choice?